

A graph theory approach to the dormitory room placement problem

Sri Efrinita Irwan¹ and Triyana Muliawati¹

¹ Mathematics Department, Institut Teknologi Sumatera, Lampung Selatan, Lampung

Abstract. One of the important areas in mathematics is graph theory. A graph is a mathematical structure used to model pairwise relations between objects. The theory of graph can be applied in various problems. The purpose of this paper is to solve the dormitory room placement problem using graph theory approach. In this paper, there are two factors used to determine the student room, i.e. the study program and provincial origin.

1. Introduction

The initial introduction of graph theory was motivated by The Seven Bridges of Königsberg problem, a historically notable problem in mathematics [1]. The city of Königsberg had seven bridges which connected two islands with the mainland via seven bridges. People staying there always wondered whether was there any way to walk over all the bridges once and only once. It was a long-standing problem until solved by Euler [2] in 1736. Euler came out with the solution in terms of graph theory. Euler pointed out that the only important feature of a route is the sequence of bridges crossed. This allowed him to reformulate the problem in abstract term. Each land mass represented by a vertex and each bridge represented by an edge. This abstraction serves to record which pair of land masses (vertices) is connected by a bridge (edge). The resulting mathematical structure is called a graph. In modern terms, a graph is known as a mathematical structure used to model pairwise relations between objects.

Graph theory is applied in various aspects such as engineering, physical, social and biological science, etc. [1]. One of the important topics in graph is graph colouring. Graph colouring has many considerable applications, such as map colouring, scheduling problem, school timetable, etc. In this paper, we discuss the application of graph colouring to the dormitory room placement problem.


Dormitory is one of the important facilities at a university. Several universities apply policies to require first-year students to live in dormitories. The existence of dormitories is often used to train student's attitude. One way that can be taken is to maintain the diversity of dormitory room residents. Example of such problems is every student in a particular room must come from a different study program and provincial origin.

2. Definitions and facts

2.1 Basic graph theory

In this subsection, we review some notions and facts related to graphs necessary for the subsequent discussion (see [3], [4], [5], and [6]). A graph is a pair $G = (V, E)$ of sets such that $E \subseteq [V]^2$; thus, the elements of E are 2-element subsets of V . We always assume that $V \cap E = \emptyset$. The elements of V are called the vertices and the elements of E are called the edges of G . The vertex set of a graph G denoted by $V(G)$, while its edge set denoted by $E(G)$.

Example 2.1 The graph H on $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ with the edge set $E = \{\{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_5\}, \{v_3, v_5\}\}$, or can be written as $E = \{e_1, e_2, e_3, e_4, e_5\}$ where $e_1 = \{v_1, v_2\}$, $e_2 = \{v_1, v_3\}$, $e_3 = \{v_1, v_4\}$, $e_4 = \{v_2, v_5\}$, $e_5 = \{v_3, v_5\}$, is drawn as follows.



Picture 1 The graph H on $V = \{v_1, v_2, v_3, v_4, v_5, v_6\}$ with the edge set $E = \{\{v_1, v_2\}, \{v_1, v_3\}, \{v_1, v_4\}, \{v_2, v_5\}, \{v_3, v_5\}\}$.

The number of vertices of a graph G is its order, denoted by $|G|$, its number of edges is denoted by $\|G\|$. Two vertices v_1, v_2 of G are adjacent, or neighbours, if $\{v_1, v_2\}$ is an edge of G , i. e. if the vertices v_1 and v_2 connected by an edge of G . The number of neighbours of a vertex v is called the degree of v .

2.2 Graph colouring

One of the important topics in graph theory is graph colouring. Graph colouring is divided into several types, including vertex colouring and edge colouring. In this paper, the graph colouring in question is a vertex colouring.


Formally, a vertex colouring of a graph $G = (V, E)$ is a map $c : V \rightarrow S$ such that $c(v_1) \neq c(v_2)$ whenever v_1 and v_2 are adjacent [3]. The elements of the set S are called the available colours. The interesting thing about S is its size. Therefore, a vertex colouring $c : V \rightarrow S$ can be restricted to $c : V \rightarrow \{1, 2, 3, \dots, k\}$ to say that G can be coloured in k different colour. The smallest positive integer k satisfying the vertex colouring rule is called a chromatic number of G , denoted by $\chi(G)$. The chromatic number in question is related to the vertex colouring of G . Henceforth, the term colouring in this paper devoted to the vertex colouring of a graph.

2.3 Welsh-Powell Algorithm

One algorithm that can be used in graph colouring is Welsh-Powell algorithm. The Welsh-Powell algorithm consists of the following step.

1. Find the degree of each vertex.
2. List the vertices in order of descending degree.
3. Colour the first vertex in the list with the colour-1.
4. Colour every vertex not adjacent to the coloured vertices using the same colour.
5. Repeat the process to the uncoloured vertices with a new colour.

Example 2.2 The chromatic number of the graph H in Picture 1 is 2.



Picture 2 A vertex colouring $c : V \rightarrow \{1, 2\}$ of H .

Based on Picture 2, it can be seen that the graph H can be coloured in two colours so that every two adjacent vertices have different colours.

3. Case Studies

Triggered by facts explained in Section 2, it is of interest to apply the theory of graph colouring to the dormitory room placement problem. Dormitory is one of the important facilities at a university. Several universities apply policies to require first-year students to live in dormitories. In managing a dormitory, a number of universities usually utilize the existence of dormitories to train student's attitude. One way that can be taken is to maintain the diversity of dormitory room residents. Example of such problems is every student in a particular room must come from a different study program and provincial origin.


Suppose in any university, given the data of fifteen first-year students who will stay in the dormitory as follows.

Table 1 The data of fifteen first-year students who will stay in the dormitory.

No	Name	Study Program	Provincial Origin
1	Student-1	Informatics Engineering	Banten
2	Student-2	Geophysical Engineering	West Java
3	Student-3	Environmental Engineering	Lampung
4	Student-4	Electrical Engineering	Lampung
5	Student-5	Environmental Engineering	Lampung

6	Student-6	Geomatics Engineering	West Sumatera
7	Student-7	Physics	North Sumatera
8	Student-8	Mathematics	North Sumatera
9	Student-9	Geophysical Engineering	North Sumatera
10	Student-10	Civil Engineering	Lampung
11	Student-11	Geomatics Engineering	Lampung
12	Student-12	Geophysical Engineering	Lampung
13	Student-13	Geomatics Engineering	Lampung
14	Student-14	Geological Engineering	Lampung
15	Student-15	Geological Engineering	South Sumatera

The problem that will be solved is how to arrange the student's room so that every student in a particular room must come from a different study program and provincial origin. Let v_i be the vertex represents the Student- i . The data in Table 1 can be represented in a graph L as follows.



Picture 3 The graph L which represents data in Table 1.

Hence it is obtained the degree of each vertex of L as in the following table.

Table 2 The degree of each vertex of L .


Vertex	v_1	v_2	v_3	v_4	v_5	v_6	v_7	v_8	v_9	v_{10}	v_{11}	v_{12}	v_{13}	v_{14}	v_{15}
Degree	0	2	7	7	7	2	2	2	4	7	8	9	8	8	1

By using Welsh-Powell algorithm, list of vertices in order of descending degree and the colour of each vertex of L is given in the following table.

Table 3 List of vertices in order of descending degree and the colour of each vertex of L .

No	Vertex	Degree	Colour
1	v_{12}	9	1
3	v_{11}	8	2
4	v_{13}	8	3
5	v_{14}	8	4
2	v_4	7	5
6	v_3	7	6
7	v_5	7	7
8	v_{10}	7	8
9	v_9	4	2
10	v_2	2	3
11	v_6	2	1
12	v_7	2	1
13	v_8	2	3
14	v_{15}	1	1
15	v_1	0	1

Therefore, the placement of student's room can be stated in the following graph.



Picture 4 The vertex colouring of L .

From the above result, we can conclude that the chromatic number of the graph L is $\chi(L) = 8$. Therefore, it can be concluded that eight rooms are needed to place fifteen students so that every student in a particular room come from a different study program and provincial origin. Note that in this case, the number of students in one room is not restricted.


The solution above is not the only solution. There are alternative solutions to this problem, including the following.

Table 4 List of vertices in order of descending degree and the colour of each vertex (an alternative).

No	Vertex	Degree	Colour
1	v_{12}	9	1
3	v_{11}	8	2
4	v_{13}	8	3
5	v_{14}	8	4
2	v_4	7	5
6	v_3	7	6
7	v_5	7	7
8	v_{10}	7	8
9	v_9	4	3

10	v_2	2	2
11	v_6	2	1
12	v_7	2	1
13	v_8	2	2
14	v_{15}	1	1
15	v_1	0	1

Therefore, the placement of student's room can be stated in the following graph.



Picture 5 The vertex colouring of L (an alternative).

Since the vertex v_1 is not adjacent to any vertex, there are 8 colouring options for the vertex- v_1 . Therefore, the Student-1 can be in the same room with any student.

4. Conclusion

Based on the explanation above, it can be concluded that for the same chromatic number, there are many alternative solutions to arrange the student's room. However, based on the existing data it is known that the majority of the students who will stay in the dormitory come from Lampung. This causes the ineffectiveness of room use.

5. References

- [1] Deo N 1990 *Graph theory with applications to engineering and computer science* (India: Prentice Hall)
- [2] Euler L 1736 Solutio problematis ad geometriam situs pertinentis *Comment. Acad. Sci. U. Petrop* **8** 128-140
- [3] Diestel R 2005 *Graph theory 3rd edition* (New York: Springer-Verlag Heidelberg)
- [4] Bapat R B 2010 *Graph and matrices* (New York: Springer London Dordrecht Heidelberg)
- [5] Bondy J A and Murty U S R 1976 *Graph Theory with Applications* (Britain: The Macmillan Press Ltd.)
- [6] Wilson R J 1996 *Introduction to graph theory 4th edition* (Longman)

Acknowledgments

The authors gratefully acknowledge the helpful assistance of Hibah Mandiri Institut Teknologi Sumatera No. 134bl/IT9.C1/PP/2018 for financial support. Also, they thank referees for their valuable comments and suggestions.